

Late Bashkirian (Early Pennsylvanian) horseshoe crabs (Chelicerata: Xiphosura) from the Mospyne Formation, Donets Basin, Ukraine

Vitaly Dernov^{1,2}

¹ Department of Palaeontology and Stratigraphy of the Palaeozoic Sediments, Institute of Geological Sciences of the NAS of Ukraine, Oles Honchar Street 55b, 01054 Kyiv, Ukraine;

² Department of Geology, National Museum of Natural History of the NAS of Ukraine, Bohdan Khmelnytskyi Street 15, 01030, Kyiv, Ukraine;

e-mail: vitalydernov@gmail.com,  <https://orcid.org/0000-0002-5873-394X>

Abstract

The Bashkirian xiphosurid assemblage of the Donets Basin comprises only three genera: *Euproops*, *Prestwichianella*, and *Belinurus*. With the exception of *Euproops danae*, all recorded species appear to be endemic to the basin. A small xiphosurid assemblage consisting of *Belinurus* sp., ?*Euproops* sp., and indeterminate xiphosurid remains is described from non-marine deposits of the Mospyne Formation (upper Bashkirian, Lower Pennsylvanian) in the central Donets Basin, eastern Ukraine. Together with *Belinurus iswarinensis* (Chernyshev, 1927), previously reported from the Bashkirian Mandrykyne and Mospyne formations, these specimens represent the oldest known Carboniferous xiphosurids from the Donets Basin.

Keywords: Xiphosura, *Belinurus*, *Euproops*, late Bashkirian, palaeoecology

1. Introduction

Extant horseshoe crabs (Xiphosura Latreille, 1802) comprise four species: *Limulus polyphemus* (Linnaeus, 1758), *Carcinoscorpius rotundicaudata* (Latreille, 1802), *Tachypleus gigas* (Müller, 1785), and *T. tridentatus* (Leach, 1819), distributed along the Atlantic coast of eastern North America and throughout southern and southeastern Asia (Lamsdell, 2019, 2020). During the Pennsylvanian, regarded as the evolutionary heyday of the group, xiphosurids were characteristic components of non-marine faunal assemblages (Selden et al., 2019; Bicknell & Pates, 2020). Their fossils are relatively common in Bashkirian to Kasimovian coal-bearing strata throughout Euramerica (e.g., Bölsche, 1885; Pack-

ard, 1886; Dix & Pringle, 1929, 1930; Pruvost, 1930; Eller, 1938; Stainier, 1938; Schwarzbach, 1962; Ambrose & Romano, 1972; Siegfried, 1972; Anderson, 1994; Schultka, 1994; Filipiak & Krawczyński, 1996; Anderson et al., 1999; Tashman et al., 2019; Lustri et al., 2020; Bicknell et al., 2019, 2021, and references therein).

In Ukraine, Pennsylvanian coal-bearing successions occur in three principal regions: the Lviv Palaeozoic Trough, the Dnipro-Donets Depression, and the Donets Basin (Shulga et al., 2007; Nemyrovska & Yefimenko, 2013; Poletaev & Vdovenko, 2013). To date, however, xiphosurids have been recorded only from the Donets Basin in eastern Ukraine (Chernyshev, 1927, 1928; Karlov, 1948; Shpinev, 2018; Dernov, 2019a, b). Their stratigraphic distribu-

tion is restricted to a relatively narrow stratigraphic interval extending from the lower Bashkirian (*Bilinguites-Cancelloceras* ammonoid genus zone) to the upper Moscovian (*Pseudoparalegoceras-Wellerites* ammonoid genus zone). This interval corresponds to the most coal-rich part of the Carboniferous succession in the basin (Logvinenko, 1953).

Chernyshev (1927, 1928) provided the first formal descriptions of Carboniferous xiphosurids from the Donets Basin, although a xiphosurid specimen had been earlier noted by Zalessky (1907). In the first paper, Chernyshev (1927) documented previously established species *Prestwichia danae* (Meek & Worthen, 1865) and described four new taxa: *Prestwichianella zalesskii*, *Bellinurus iswarinensis*, *B. metschetsnensis*, and *B. stepanowi*, all from the Pennsylvanian strata of the basin. Subsequently, Chernyshev (1928) briefly described and illustrated new specimens of *Prestwichia danae*, *Bellinurus iswarinensis*, and *B. stepanowi*. Because the generic name *Prestwichia* Woodward, 1867 is preoccupied, it was replaced by *Prestwichianella* Cockerell, 1905 (Shpinev, 2018).

Karlov (1948) later described a further species, *Prestwichianella tschernyshewi*, but its taxonomic validity remains uncertain. Shpinev (2018) partly revised the taxa established by Chernyshev and examined additional material from the Donets Basin. Shpinev (2018) and Bicknell & Pates (2020) illustrated several specimens of the type series of the species described by Chernyshev (1927). Nevertheless, the species of *Bellinurus* erected by Chernyshev (1927) require comprehensive revision to clarify their generic assignments (Lamsdell, 2022).

The present study formally documents new xiphosurid material from the upper Bashkirian Mospyne Formation of the Donets Basin. These fossils represent some of the oldest horseshoe crabs known from the Carboniferous succession of the basin. Previously, Chernyshev (1928) briefly described and figured *Bellinurus iswarinensis* from the roof shale of the g_3 coal in the Mospyne Formation. This formation has yielded a diverse arthropod fauna, including trilobites (Weber, 1933; Mychko & Dernov, 2019), arthropleurid myriapods (Dernov, 2019a, 2025b), conchostracans (Dernov & Udovychenko, 2021), cyclids (Dernov, 2022c), phyllocarids (Dernov, 2025a), and insects (Dernov, 2019a, 2019b, 2025b). Xiphosurid remains from the Mospyne Formation have also been briefly mentioned by the present author in earlier studies (Dernov, 2019a, 2019b, 2025a).

The aims of this study are twofold: (1) to document the systematic diversity of horseshoe crabs from the early late Bashkirian of the Donets Basin,

and (2) to discuss their palaeoenvironment, habitat preferences, and taphonomic conditions.

2. Materials and methods

Twelve poorly to moderately preserved impressions and compressions of xiphosurid exoskeletal fragments were examined. Owing to the generally poor preservation of the material, the systematic placement of a substantial proportion of the material remains uncertain.

The material described herein was collected by the author from three stratigraphic horizons within the Mospyne Formation (Zuyivkian Horizon, lower part of the Kayalian regional stage, upper Bashkirian, Lower Pennsylvanian; Fig. 1A) in southern Luhansk Oblast, Ukraine. These horizons comprise: (1) a 0.7 m thick interval of greyish-yellow shale interbedded within a deltaic sandstone succession, approximately 43 m below the G_1^2 limestone; (2) the roof shale of the g_1^2 coal seam; and (3) the roof shale of the g_2 coal seam (Fig. 1B).

The fossil-bearing horizons are exposed at three localities in the vicinity of the village of Makedonivka, Luhansk Raion, Luhansk Oblast (Fig. 2): (1) sandstone quarries approximately 1 km west of the village (48°14'12.0"N, 39°15'19.4"E); (2) spoil heaps of an abandoned small coal mine near the northern




Fig. 1. A – Stratigraphic position of the Mospyne Formation within the Carboniferous succession of the Donets Basin. B – Stratigraphic position of the xiphosurid-bearing horizons within the Mospyne Formation. Abbreviations: Fm, Formation; Gr, Group; M., Mokra.

outskirts of the village, where the g_1^2 coal seam was formerly exploited (48°14'36"N, 39°17'58"E); and (3) spoil heaps of small-scale coal mines approximately 1.5 km north of the village, where the g_2 coal seam was mined (48°14'54.1"N, 39°18'20.6"E).

Five impressions and compressions of prosomal and thoracetrionic remains assigned to *?Euproops* sp. and *Xiphosurida* indet. (specimens NMNHU G-10659/01-05) were recovered from Horizon 1 (Fig. 3A). These are preserved within small weathered siderite concretions embedded in shale associated with the deltaic sandstone succession.

Horizon 2 (Fig. 3B, C) yielded three impressions and compressions of thoracetrionic fragments referred to *Belinurus* sp. and *Xiphosurida* indet. (GMLNU-56/06-08). These specimens occur in black carbonaceous shale and in siderite and limonite concretions derived from several beds within the roof-shale succession above the g_1^2 coal seam. The best-preserved specimen (GMLNU-56/06) was collected from greyish-yellow siltstone exposed immediately above the entrance to the abandoned mine.

Three specimens consisting of the prosomal and thoracetrionic remains assigned to *?Euproops* sp. and *Xiphosurida* indet. (GMLNU-56/09-11) were collected from Horizon 3 (Fig. 3D). They are preserved in siderite and limonite concretions enclosed within the roof shale of the g_2 coal seam.

Most specimens are preserved either within siderite and limonite concretions or as impressions on shale bedding planes. The concretions are strongly weathered as a result of prolonged exposure on spoil heaps or shallow burial in the bedrock. The original preservation of the xiphosurid remains was probably considerably better, but weathering of the concretions substantially reduced the quality of preservation. In some cases (e.g. Fig. 6F), weathering of the concretion core produced hollow cavities containing detached internal moulds of the exoskeleton that move freely within the concretion. Such heavily weathered concretions are known locally as "rattles" (*kolotushky* in Ukrainian; Logvinenko, 1953). Most xiphosurid remains occur as flattened exoskeletal fragments preserved parallel to the bedding plane, although several specimens




Fig. 2. Localities yielding horseshoe crabs. A – Geological map of the southern part of Luhansk Oblast (modified after Dernov, 2025a). B – Detailed map of the area outlined in Fig. 2A.



Fig. 3. General views of the xiphosurid-bearing localities (A–D) and biota associated with horseshoe crabs from fossiliferous Horizon 1 (E–M). **A** – Siltstone exposure at fossil site No. 1; **B** – Spoil heap of an abandoned coal mine at fossil site No. 2; **C** – Slope above the entrance to an abandoned coal mine at fossil site No. 2; **D** – Spoil heap of an abandoned coal mine at fossil site No. 3; **E–J** – Non-marine bivalves from fossiliferous Horizon 1: **E** – *Carbonicola rectilinearis* Trueman & Weir, 1948 (IGS NASU-11/03); **F** – *Carbonicola limax* Wright, 1934 (IGS NASU-11/15); **G–J** – Shells showing injuries sustained during life (**G**, **H**) and blister pearl-like structures (**I**, **J**) (**G** – IGS NASU-11/09; **H** – IGS NASU-11/14; **I** – IGS NASU-11/17; **J** – IGS NASU-11/16); **K** – Poorly preserved prosoma of a horseshoe crab from the sandstone of fossiliferous Horizon 1 (GMLNU-56/01); **L** – Poorly preserved pygocephalomorph crustacean from fossiliferous Horizon 1 (field photograph); **M** – Evidence of a possible fungal-plant interaction on a lycopsid leaf (field photograph). Scale bars = 5 mm.

(e.g. Fig. 6D, G) appear to have been preserved in an enrolled posture.

One uncatalogued specimen illustrated in Fig. 3K is preserved in sandstone and was probably transported prior to burial, as indicated by its poor preservation and its association with fragments of arthropleurid exoskeletons and plant stems (Dernov, 2019a). In contrast, the remaining specimens are interpreted as autochthonous or parautochthonous because neither the xiphosurid remains nor the associated faunal elements exhibit evidence of transport.

Although the studied material originates from only three stratigraphic horizons, it represents the outcome of many years of field collecting. Non-marine deposits within the Mospyne Formation, particularly the lacustrine roof shales of the coal seams that have yielded most Carboniferous xiphosurids from the Donets Basin (Chernyshev, 1927, 1928; Karlov, 1948; Shpinev, 2018), are uncommon in this formation. By contrast, the overlying Smolyanynivka Formation, deposited during the late Bashkirian regression (Ustinovsky, 1993), contains considerably more extensive non-marine facies and preserves a much richer non-marine biota (Logvinenko, 1953).

The studied xiphosurid specimens are housed in the Department of Geology, National Museum of Natural History, National Academy of Sciences of Ukraine, Kyiv (NMNHU G-10659), and the Geological Museum of the Taras Shevchenko National University of Luhansk, Poltava (GMLNU-56). Collections of associated fossils are deposited in the Department of Geology, National Museum of Natural History, National Academy of Sciences of Ukraine, Kyiv (NMNHU-G 8640), the Department of Stratigraphy and Palaeontology of the Palaeozoic Sediments, Institute of Geological Sciences, National Academy of Sciences of Ukraine, Kyiv (IGS NASU-11, IGS NASU-15, IGS NASU-41, and two uncatalogued specimens), and the Geological Museum of Taras Shevchenko National University of Luhansk, Poltava (GMLNU-9 and GMLNU-19).

Morphological terminology and measurement protocols follow Filipiak & Krawczyński (1996). The systematic framework adopted for the family Belinuridae von Zittel in von Zittel & Eastman, 1913, including the spelling of the type genus, follows Lamsdell (2020) and Lamsdell & Clapham (2021).

3. Geological setting

The Donets Basin forms the southeastern segment of the northwest-southeast-trending Pripyat-Dni-pro-Donets intracratonic rift system, which extends

from the Baltic region to the Caspian Sea across Belarus, Ukraine, and Russia. The rift is situated between the Voronezh Massif to the north and the Ukrainian Shield to the south and occupies the southwestern margin of the East European Platform (Izart et al., 1996; Eros et al., 2012; Sachsenhofer et al., 2012; Van Hinsbergen et al., 2015).

The Mospyne Formation is a 315–730 m thick cyclic succession composed of mudstones, siltstones, sandstones, limestones, and coal seams (Aisenverg et al., 1963, 1975; Feofilova & Levenstein, 1963; Dunaeva, 1969; Poletaev et al., 2011; Nemyrovska & Yefimenko, 2013). In the study area, the lower part of the formation (below the g_1 coal seam) is replaced by a shale-dominated succession that is notably poor in fossils. This interval is most likely assignable to the Dyakove Group (upper Visean–upper Bashkirian), a predominately argillaceous unit recognized throughout the southern Donets Basin (Reznikov, 1993; Dernov, 2022b; Dernov & Poletaev, 2024).

The Bashkirian coal-bearing succession accumulated predominantly on an extensive alluvial-deltaic plain along the southeastern margin of Laurussia. Deposition took place under alternating continental and marginal-marine conditions, with repeated incursions from the epicontinental sea producing the characteristic cyclothemic succession of the basin (Logvinenko, 1953; Feofilova & Levenshtein, 1963; Ustinovsky, 1993). During the Bashkirian, marine sedimentation was concentrated mainly in the central part of the Donets Basin, whereas marginal areas, including the study area, remained dominated by continental depositional environments (Reznikov, 1993; Dernov & Poletaev, 2024).

4. Associated biota

Macroscopic plant remains assigned to genera *Cyperites*, *Lepidodendron*, *Stigmaria*, *Asterophyllites*, *Eusphenopteris*, *Karinopteris*, and *Cordaites*, as well as non-marine bivalves *Carbonicola rectilinearis* Truman & Weir, 1948 (Fig. 3E) and *C. limax* Wright, 1934 (Fig. 3F), and a poorly preserved impression of a pygocephalomorph crustacean caparace (Fig. 3L), co-occur with xiphosurids in fossiliferous Horizon 1. Some shells of *Carbonicola* exhibit injuries sustained during life (Fig. 3G–J) (Dernov, 2022d). Lycopsid leaves are commonly preserved in 3D form owing to carbonate mineralisation and occasionally show evidence of fungal colonisation (Fig. 3M).

Xiphosurid resting traces (*Cubichnia*) assigned to the ichnogenera *Arborichnus*, *Selenichnites*, and *Crescentichnus*, occur at several stratigraphic lev-




Fig. 4. Biota associated with horseshoe crabs from fossiliferous Horizon 2. **A, B** – ?Paratergites of the myriapod *Arthropleura* sp. (**A** – NMNHU-G 8640/01; **B** – GMLNU-19/01); **C** – Pteridosperm *Neuraethopteris schlehanii* (Stur) Cremer (uncatalogued); **D** – Pteridosperm *Paripteris gigantea* (Sternberg) Gothan (uncatalogued); **E** – Galls on a plant stem (uncatalogued); **F, G** – Bivalves *Curvimirula trapeziforma* (Dewar, 1939) (**F** – IGS NASU-11/32; **G** – IGS NASU-11/34); **H** – Bivalve *Curvimirula tessellata* (Jones, 1891) (IGS NASU-11/43); **I** – Gastric residue composed of fragmented fish scales (uncatalogued); **J** – Limonitised coprolite (uncatalogued); **K** – Trace fossil *Cochlichnus anguineus* Hitchcock, 1858 (GMLNU-9/02); **L** – Fish scale belonging to *Rhizodopsis sauroides* (Williamson, 1849) (uncatalogued). Scale bars = 5 mm.

els within the Mospyne Formation, including the sandstone associated with fossiliferous Horizon 1 and the sandstone overlying fossiliferous Horizon 2 (Dernov, 2019a, 2023b). All these trace fossils originate from deltaic and prodeltaic deposits, suggesting that their producers were likely marine xiphosurids.

The width of *Crescentichnus* resting traces from the sandstone associated with fossiliferous Horizon 1 indicates that the prosoma of some trace-making xiphosurids measured approximately 15 cm across (Dernov, 2019a). If so, the producers of these traces this would rank among the largest Carboniferous xiphosurids known. At present, the largest known Carboniferous horseshoe crab is *Xaniopyramis linseyi* Siveter & Selden, 1987, from the Serpukhovian of England, with a prosomal width of approximately 14 cm (Siveter & Selden, 1987). *Crescentichnus* is associated with arthropleurid exoskeleton fragments, plant stem remains, and pebbles (Dernov, 2019a). A poorly preserved compression of the prosoma of an unidentified xiphosurid was also discovered at this level (Fig. 3K).

The roof shale of the g_1^2 coal seam (fossiliferous Horizon 2) yields a diverse fossil assemblage comprising the non-marine bivalves *Curvirimula trapeziforma* (Dewar, 1939) (Fig. 4F, G) and *C. tes-*

selata (Jones, 1891) (Fig. 4H); the myriapod *Arthropleura* sp. (Fig. 4A, B); scales of crossopterygian fishes including *Rhizodopsis sauroides* (Williamson, 1849) (Fig. 4L) and *Rhabdoderma elegans* (Newberry, 1856); a bradyodont tooth referred to *Helodus* sp.; and fragmentary fish skeletal remains. Also present are macroproblematica, trace fossils including burrows *?Palaeophycus* isp., locomotion traces such as *Cochlichnus anguineus* Hitchcock, 1858 (Fig. 4K) and *Diplichnites* isp., bromalites (Fig. 4I, J), and evidence of plant-arthropod interactions, including galling, endophytic oviposition, and margin feeding (Fig. 4E) (Dernov, 2019b, 2021, 2022a, 2023a, 2025b).

The best-preserved specimen (Fig. 3A), assigned to *Belinurus* sp., was recovered directly from the outcrop face, in contrast to all other specimens, which were collected from the spoil heap. Associated with this specimen were impressions of the terrestrial plants *Calamites* and *Karinopteris*, bivalves, and trace fossils *?Palaeophycus* isp. and *Cochlichnus anguineus* (Dernov, 2019b; 2023a).

A diverse assemblage of terrestrial plants was also collected from fossiliferous Horizon 2, including the genera *Asolanus*, *Bothrodendron*, *Cyperites*, *Halonia*, *Lepidodendron*, *Lepidophloios*, *Lepidostrobo-*




Fig. 5. Biota associated with horseshoe crabs from fossiliferous Horizon 3. A – Bivalve *Naiadites* sp. (IGS NASU-11/45); B – Bivalve *Carbonicola obtusa* (Hind, 1894) (IGS NASU-11/21); C – Phyllocarid crustacean *Dithyrocaris* cf. *granulata* Woodward & Etheridge, 1873 (IGS NASU-41/01); D – Cyclidan crustacean *Brittaniclus* cf. *scotti* (Woodward, 1893) (IGS NASU-15/02); E, H – Insect wings (uncatalogued); F – Conchostracan *Pseudestheria* sp. (uncatalogued); G – Indeterminate conchostracan crustacean (uncatalogued). Scale bars = 5 mm.

eris, *Dictyoxylon*, *Eusphenopteris*, *Neuraethopteris* (Fig. 4C), and *Paripteris* (Fig. 4D) (Dernov & Udovychenko, 2019).

An assemblage of terrestrial plant remains has been recovered from the roof shale of the g₂ coal seam (fossiliferous Horizon 3), including *Bothrodendron*, *Cyperites*, *Lepidodendron*, *Lepidophloios*, *Sigillaria*, *Stigmaria*, *Syringodendron*, *Calamites*, *Pinnularia*, *Dictyoxylon*, *Karinopteris*, *Paripteris*, *Trigonocarpus*, *Cordaianthus*, *Cordaicarpus*, and *Cordaites* (Dernov, 2025a). In addition to plants, the site has yielded a variety of non-marine invertebrates, including microconchids; the bivalves *Carbonicola obtusa* (Hind, 1894) (Fig. 5B), *Curvirimula trapeziforma* (Dewar, 1939), and *Naiadites* sp. (Fig. 5A); the conchostracans *Pseudestheria* sp. (Fig. 5F); and the phyllocarid crustacean *Dithyrocaris* cf. *granulata* Woodward & Etheridge, 1873 (Fig. 5C). Other notable finds include a fragmentary impression of a wing base of a large insect (Fig. 5H), plant remains showing arthropod-induced damages, and pseudofossils assigned to *Astropolithon* (Dernov, 2021, 2022a, 2025a). The cyclid crustacean *Brittaniclus* cf. *scotti* (Woodward, 1893), previously reported from the base of the g₂ coal seam (Dernov, 2022c) (Fig. 5D), may in fact have originated from the roof shale of this coal seam. It was found in association with an assemblage dominated by terrestrial plants, as well as myriapods, conchostracans (Fig. 6G), and insects (Fig. 6E).

5. Age of the xiphosurid fauna

The Mospyne Formation contains numerous biostratigraphic indicators that constrain its age to the early late Bashkirian (equivalent to the Langsettian or Westphalian A of the West European stratigraphic scale). These include characteristic terrestrial plant assemblages containing stratigraphically significant species *Sigillaria elegans* Brongniart, *Lyginopteris hoeninghausi* Potonié, *Neuraethopteris rectinervis* (Kidston) Laveine, *N. schlehanii* (Stur) Cremer, and *Karinopteris acuta* (Brongniart) Boersma (Novik, 1974; Fissunencko, 1991; Boyarina, 2016; Dernov & Udovychenko, 2019); ammonoids of the *Branneroceras*-*Gastrioceras* Genus Zone (Popov, 1979; Dernov, 2022b); and non-marine bivalves belonging to *Curvirimula trapeziforma* (Dewar, 1939), *C. tessellata* (Jones, 1891), *Carbonicola rectilinearis* Trueman & Weir, 1948, *C. limax* Wright, 1934, which indicate the upper part of the *lenisulcata* Zone and the lower part of the *communis* Zone (Dernov, 2022a).

Conodonts identified from these strata include ‘*Streptognathodus*’ *expansus* Igo & Koike, 1964, ‘*S.*’ *su-*

berectus Dunn, 1966, *Idiognathoides sinuatus* Harris & Hollingsworth, 1933, *I. corrugatus* Harris & Hollingsworth, 1933, *I. sulcatus* Higgins & Bouckaert, 1968, and *Idiognathodus sinuosus* Ellison & Graves, 1941, all of which indicate the ‘*Streptognathodus*’ *expansus* Zone (Kozitskaya et al., 1978; Nemyrovskaya, 1999, 2017). This zone is correlated with the Langsettian *Idiognathoides sulcatus parvus* Zone of England, the *Idiognathodus klapperi* Zone of the upper Morrowan (Pennsylvanian) of the United States, and the lower part of the *Declinognathodus marginodosus* Zone in the historical stratotype of the Bashkirian Stage in Bashkiria, Russia (Nemyrovskaya, 1999).

6. Systematic palaeontology

The best-preserved specimens of horseshoe crabs, assigned to *Belinurus* sp. and ?*Euproops* sp., are described below. The more poorly preserved fossils are only illustrated (Fig. 6F–I).

Order Xiphosurida Latreille, 1802

Suborder Belinurina von Zittel in von Zittel & Eastman, 1913

Family Belinuridae von Zittel in von Zittel & Eastman, 1913

Genus *Belinurus* Bronn, 1839

Type species. *Limulus trilobitoides* Buckland, 1837; by subsequent designation.

Included species. *Belinurus carwayensis* Dix & Pringle, 1929; *B. concinnus* Dix & Pringle, 1929; *B. grandaevus* Jones & Woodward, 1899; *B. kiltorkensis* Baily, 1869; *B. morgani* Dix & Pringle, 1930; *B. pustulosus* Dix & Pringle, 1929; *B. silesiacus* (Roemer, 1883); *B. sustai* (Prantl & Přibyl, 1956); *B. trechmanni* Woodward, 1918 (Lamsdell & Clapham, 2021).

Diagnosis. Belinurid with axis of first thoracetrone tergite medially inflated; thoracetrone ovoid to semicircular in outline; thoracetrone fixed tergo-pleural spines elongate, needle-like (after Lamsdell & Clapham, 2021, p. 1354).

Remarks. The correct authorship and spelling of *Belinurus* was recently clarified by Lamsdell & Clapham (2021), who demonstrated that *Belinurus* Bronn, 1839 was the correct spelling and attribution as opposed to *Bellinurus* Pictet, 1846.

Belinurus sp.

Fig. 6A

Material. One specimen (GMLNU-56/06) from fossiliferous Horizon 2 (fossil site No. 2).

Description. Specimen GMLNU-56/06 is preserved as a compression of the dorsal surface of the thoracetron and the proximal part of the telson within a siderite nodule. The thoracetron measures 9.5 mm in length and 15 mm in width at its broadest point (width/length ratio = 1.58). It is ovoid to weakly subtriangular in outline and consists of eight segments, each 0.6–1.0 mm wide. The thoracetronic axis is 4.2 mm wide anteriorly, where it reaches its maximum width, and tapers gradually posteriorly. The axial segments and the microtergite are not preserved. The lateral ends of

the thoracetronic segments bear poorly preserved, narrow spines that are gently recurved posteriorly. The best-preserved spine, on the fourth segment of the right side, is 3 mm long. A roll-shaped callosity extends along the posterior margin of each segment from the pleural region to the base of the corresponding spine. The thoracetronic flange is narrow, approximately 0.5 mm wide, and maintains a nearly constant width throughout. Only the proximal part of the telson is preserved; it is approximately 12 mm long and 1 mm wide at the base and bears a distinct median ridge.




Fig. 6. Xiphosurids from the Mospyne Formation. **A** – *Belinurus* sp. (compression of the thoracetron in dorsal view, GMLNU-56/06); **B–E** – *Euproops* sp. (**B** – impression of the prosoma and thoracetron in dorsal view, NMNHU G-10659/01; **C** – compression of the thoracetron with part of the prosoma, in dorsal view, GMLNU-56/09; **D** – enrolled exoskeleton, NMNHU G-10659/04; **E** – impression of a genal spine, NMNHU G-10659/03); **F–I** – Xiphosurida indet. (**F, G** – enrolled exoskeletons, GMLNU-56/11 and GMLNU-56/08; **H** – impression of the prosoma in dorsal view, NMNHU G-10659/02; **I** – compression of the prosoma with a partially preserved thoracetron, in dorsal view, GMLNU-56/10). Scale bars = 5 mm.

Remarks. The specimen described above is referred to *Belinurus* on the basis of its ovoid thoracetrone bearing elongate, needle-like fixed tergopleural spines. It differs from *Belinurus iswarinensis*, specimens of which have been reported from several localities, including the Mospyne Formation (Chernyshev, 1927, 1928), in possessing (1) a weakly subtriangular thoracetrone rather than the semicircular outline characteristic of *B. iswarinensis* and (2) considerably more slender spines at the lateral margins of the thoracetrone segments. On the basis of the overall thoracetrone outline and the presence of long, slender pleural spines, the specimen described here most closely resembles *Belinurus metschetsnensis* (cf. the specimen illustrated in Shpinev, 2018, fig. 3B, and Bicknell & Pates, 2020, fig. 15D). However, the poor preservation of the present specimen precludes a detailed morphological comparison with other representatives of the genus *Belinurus*.

Genus *Euproops* Meek, 1867

Type species. *Bellinurus danae* Meek & Worthen, 1865; by original designation.

Included species. Type species only.

Diagnosis. Belinurid characterised by ophthalmic ridges that curve axially posterior to the lateral eyes; a cardiac lobe bordered by dorsal furrows; elongate ophthalmic spines arising from the posterior ends of the ophthalmic ridges and extending over the thoracetrone; proximally expanded fixed tergopleural spines forming an opisthosomal flange; and a conical opisthosomal boss (after Lamsdell, 2020, p. 14).

?*Euproops* sp.

Fig. 6B–E

Material. Four specimens (NMNHUG-10659/01, NMNHU G-10659/03, NMNHU G-10659/04, and GMLNU-56/09) from fossiliferous horizons 1 and 3 (fossil sites Nos. 1 and 3).

Description. Specimen NMNHU G-10659/01 (Fig. 6B) is the best-preserved specimen referred to ?*Euproops* sp. It consists of a weathered, fragmentary impression of the prosoma and thoracetrone preserved within a siderite nodule. The prosoma is semicircular in outline, measuring approximately 11 mm in width and 3.8 mm in length. A single, slender genal spine is preserved; it is 2.5 mm long, 0.7 mm wide at the base, and oriented subparallel to the longitudinal axis of the exoskeleton. The cardiac lobe is small, subtriangular, and tapers anteriorly. The paired ophthalmic ridges curve axially posterior to the small lateral compound eyes. The median

ocelli are situated anteriorly at the point where the ophthalmic ridges converge. The thoracetrone measures 6.5 mm in length and 8.5 mm in maximum width (width/length ratio = 1.31). It is fully fused and has a semicircular to weakly subtriangular outline. The pleural segments and the axial structures of the thoracetrone are not preserved. The thoracetrone axis is narrow (approximately 1.5 mm wide), but its outline cannot be determined.

Specimen GMLNU-56/09 (Fig. 6C) is preserved as an impression of the dorsal surface of the thoracetrone together with the posterior part of the prosoma within a siderite nodule. The prosoma is very poorly preserved, and its morphology cannot be determined. The thoracetrone measures approximately 8 mm in length and 13 mm in maximum width (width/length ratio c. 1.63), and has a subtriangular to nearly semicircular outline. It appears to consist of seven segments, each 0.5–0.9 mm long. The thoracetrone axis is cylindrical and approximately 4.0 mm wide at its widest point. The axial segments and microtergite are poorly preserved. The lateral and posterior margins of the thoracetrone bear a poorly preserved flange with narrow spines that curve slightly posteriorly and are approximately 2.0–2.5 mm long. A ridge-like callosity extends along the posterior margin of each segment from the pleural region to the base of the corresponding spine. Only the proximal part of the telson is preserved, with a basal width of approximately 2 mm.

Remarks. An additional specimen (Fig. 6D), representing an enrolled horseshoe crab, probably also belongs to ?*Euproops* sp. However, owing to its poor state of preservation, this assignment remains tentative. Specimen NMNHU G-10659/03 (Fig. 6E) consists of an impression of the lateral part of the prosoma bearing a slender, elongate genal spine.

The poor preservation of the available material precludes confident assignment to *Euproops*, as it does not permit determination of the number of opisthosomal segments, assessment of the morphology of the fixed tergopleural spines, or confirmation of the presence of ophthalmic spines. Nevertheless, the specimens are tentatively referred to ?*Euproops* on the basis of several morphological features, including the course of the ophthalmic ridges, the relatively slender genal spines oriented nearly parallel to the longitudinal axis of the exoskeleton, the semicircular prosoma, the small subtriangular cardiac lobe, and the position of the median ocelli.

7. Discussion

Palaeoenvironments. According to the available lithological and palaeontological evidence, fossiliferous Horizon 1 was deposited in a shallow, low-energy lagoon with reduced salinity or in a freshwater lake (Dernov, 2022a). The bivalves from fossiliferous Horizon 2 are interpreted to have inhabited shallow, freshwater or brackish lakes situated on a coastal alluvial plain. These water bodies were characterised by low biodiversity, slow sedimentation rates, dysaerobic bottom-water conditions, and a continuous supply of nutrients from the surrounding wetlands (Dernov, 2022a, 2025b).

Bivalves of the genus *Curvirimula* probably adopted a pseudoplanktonic mode of life, attaching themselves to floating substrates such as fragments of wood or to benthic organisms, including macroscopic algae (Warth, 1967; Wignall & Sims, 1990). This life habit enabled them to remain above the oxygen-depleted bottom waters. However, the pseudoplanktonic habit was probably facultative, and under more favourable environmental conditions these bivalves may also have adopted a benthic mode of life (Dernov, 2022a). *Curvirimula* is well known from brackish-water depositional environments (Betekhtina, 1974; Eagar, 1981; Anderson et al., 1997).

The margins of these lakes were occupied by dense but taxonomically monotonous hydrophilic vegetation, dominated by monospecific stands of the arborescent sphenopsid *Calamites* (Fissunenkov, 1987).

The fossils from fossiliferous horizons 2 and 3 were collected from mine spoil heaps and most probably originated from different parts of the roof shales of the g_1^2 and g_2 coal seams. In the Donets Basin, the roof shales of coal seams typically comprise several palaeoenvironmental zones (facies) representing different depositional settings (Chernyshev, 1931; Yefimov, 1934; Logvinenko, 1953; Posudiyevsky, 1977; Dernov, 2022a): (1) swamp (lower part) to lacustrine (upper part) facies, represented by black mudstones and siltstones containing abundant plant debris, non-marine bivalves, and conchostracans; (2) brackish to shallow-marine facies, represented by black mudstones and siltstones with lingulide brachiopods, marine bivalves, and ostracods; and (3) offshore marine facies, represented by shales containing marine bivalves, gastropods, ammonoids, and other typical marine fauna. The horseshoe crabs described here most probably originated from the lacustrine subdivision of facies zone 1 (Dernov, 2025a).

These lakes developed on an alluvial-deltaic swampy plain (Zhemchuzhnikov et al., 1960; Feofilova & Levenshtein, 1963; Zaritskiy, 1970; Dernov, 2022a, 2025a). Their occurrence directly above the coal seams indicates that they formed following the burial of peat mires by clastic sediments and the consequent cessation of peat accumulation. The lake-bottom muds were probably enriched in hydrogen sulphide as a result of sulphate-reducing bacterial activity, stimulated by the abundant organic detritus, chiefly fragments of terrestrial plants transported into the lakes from the surrounding highly productive wetlands (Dernov, 2022a, 2025a). Hydrogen sulphide probably accumulated periodically within the water column or may even have persistently contaminated its lowermost part (Dernov, 2022a, 2025a).

General characteristics of the xiphosurid faunule. No xiphosurids have previously been described or illustrated from the study area, namely the central Donets Basin. However, Shpinev (2018: pl. 10, fig. 6; text-fig. 6) figured and described a juvenile specimen of *Belinurus* sp. from an abandoned coal quarry located approximately 3.3 km south of the village of Zakhidne, Luhansk Raion, Luhansk Oblast, about 23 km northwest of the village of Makedonivka. The specimen was collected from the upper Bashkirian Smolyanivka Formation, probably from shale beds associated with the h_{10} coal seam near the top of the formation. Biostratigraphically, these strata belong to the terminal *Branneroceras*–*Gastrioceras* Genus Zone or the lowermost *Diaboloceras*–*Axinolobus* Genus Zone (Popov, 1979; Dernov, 2022b, e).

The juvenile specimen figured by Shpinev closely resembles the *Belinurus* sp. described herein in possessing long, slender pleural spines. However, it differs in several respects. Its thoracetrone is semicircular in outline, whereas that of *Belinurus* sp. described herein is weakly subtriangular, and its prosoma is proportionally much broader than the thoracetrone.

The horseshoe crabs described herein, together with *Belinurus iswarinensis* described by Chernyshev (1927, 1928) from the roof shale of the f_1 coal in the lower Bashkirian Mandrykyne Formation and from the roof shale of the g_3 coal in the Mospyne Formation, currently represent the oldest known Carboniferous xiphosurids from the Donets Basin. In total, the following xiphosurid species have been reported from the Bashkirian of the Donets Basin: *Euproops danae*, ?*Euproops* sp., *Prestwichianella tschernyshevi*, *Belinurus iswarinensis*, *B. metschetsnensis*, and *B.* sp. (Chernyshev, 1927, 1928; Karlov, 1948; Shpinev, 2018; this study).

System	Subsystem	Stage	Lithostratigraphic unit	Stratigraphic range of xiphosurids	Ammonoid genus zones (for Pennsylvanian part of the succession)	Conodont zones (for Pennsylvanian part of the succession)
CARBONIFEROUS	PENNSYLVANIAN	Gzhelian	Kartamysh Fm.	<i>B. isvarinensis</i> <i>B. metschelnensis</i> <i>B. stepanowi</i> <i>B. spp.</i> <i>E. danae</i> <i>?E. sp.</i>	Not zoned	Not showed
			Araukarytova Fm.			
		Kasim.	Avilovka Fm.			<i>Swadelina subexcelsa</i>
			Isayivka Fm.			<i>Swadelina gurkovaensis</i> + <i>Swadelina dissecta</i>
		Moscovian	Gorlivka Fm.			<i>Idiogonathodus izwaricus</i>
			Almazna Fm.			<i>"Streptognathodus" transitivus</i>
			Kamensk Fm.			<i>Neognathodus atokaensis</i>
		Bashkirian	Belaya Kalitva Fm.			<i>Decl. donetzianus</i>
			Smolyanyivka Fm.			<i>Decl. marginodosus</i>
			Mospyne Fm.			<i>Idiogonathoides tuberculatus</i> - <i>Id. fossatus</i>
			Mandrykyne Fm.			<i>"Streptognathodus" expansus</i>
			Amvrosiyivka Fm.			<i>Idiogonathodus sinuosus</i>
	MISSISSIPPIAN	T. Visean/Serpukhovian	Kalmius Fm.			<i>Idiogonathoides sinuatus</i> - <i>Id. sulcatus sulcatus</i>
			Samara Fm.			<i>Declinognathodus noduliferus</i>
			Mezha Fm.			Not showed
			Mokra Volnovakha Group			

Fig. 7. Stratigraphic distribution of horseshoe crab taxa within the Carboniferous succession of the Donets Basin (after Chernyshev, 1927, 1928, 1939; Shpinev, 2018; and this study). Abbreviations: B., *Belinurus*; D., *Declinognathodus*; E., *Euproops*. Ammonoid zonation after Popov (1979) and Dernov (2022e); conodont zonation after Nemyrovska (2017). Note: The precise stratigraphic position of the locality yielding the type series of *Prestwichianella tschernyshevi* Karlov, 1948, is unknown.

At present, the Bashkirian xiphosurid assemblage of the Donets Basin comprises only three genera: *Euproops*, *Prestwichianella*, and *Belinurus* (Fig. 7). Almost all recorded species appear to be endemic to the basin, with the exception of *Euproops danae*, which is widely distributed in Pennsylvanian deposits throughout Euramerica (e.g. Meek, 1867; Fisher, 1977, 1979; Brauckmann, 1982; Anderson, 1994; Mikulic, 1998; Babcock & Merriam, 2000; Haug et al., 2012; Lucas et al., 2014; Tashman et al., 2019; Bicknell et al., 2021, and references therein).

The generally poor preservation of xiphosurids in the Mospyne Formation limits detailed comparisons with other palaeoequatorial xiphosurid faunas of Euramerica. In contrast, other arthropod groups from the Mospyne Formation, including phyllocarids and cyclids identified to species level (Dernov, 2022c, 2025a), exhibit much broader palaeobiogeographical distributions and are also known from Westphalian (upper Bashkirian–Moscovian) strata of the United Kingdom.

7. Conclusions

The xiphosurid fauna of the Mospyne Formation provides new insights into the diversity, distribution, and palaeoecology of Carboniferous horseshoe crabs in the Donets Basin. New material recovered from three fossiliferous horizons documents the presence of at least two belinurid taxa, referred to *Belinurus* sp. and *?Euproops* sp., among the oldest known Carboniferous xiphosurids from the basin.

The specimens were recovered from deposits formed in low-energy continental to marginal environments, including freshwater and brackish lakes developed on an alluvial–deltaic swampy plain. The associated faunal and floral assemblages, together with sedimentological evidence, indicate that these horseshoe crabs inhabited shallow water bodies characterised by abundant organic input and periodically dysaerobic bottom waters. These environmental conditions are consistent with the ecological preferences inferred for Pennsylvanian belinurids elsewhere in Euramerica.

Biostratigraphic evidence derived from terrestrial plants, ammonoids, non-marine bivalves,

and conodonts constrains the age of the xiphosurid-bearing strata to the early late Bashkirian (equivalent of the Langsettian of the Western European chronostratigraphic scheme). The occurrence of *Belinurus* and ?*Euproops* in these deposits extends the documented record of Carboniferous horseshoe crabs in eastern Euramerica.

The Bashkirian xiphosurid assemblage of the Donets Basin currently comprises three genera – *Belinurus*, *Euproops*, and *Prestwichianella* – with most species apparently endemic to the basin. Although the available material is generally too poorly preserved to permit detailed taxonomic and palaeobiogeographical comparisons, the new discoveries demonstrate that horseshoe crabs were a regular component of Bashkirian non-marine and marginal-water ecosystems in the Donets Basin and provide an important contribution to our understanding of the early evolutionary and palaeobiogeographical history of Carboniferous xiphosurids.

Acknowledgements

The author would like to thank to Dr Jason A. Dunlop (Museum für Naturkunde – Leibniz Institute for Research on Evolution and Biodiversity, Berlin) and Dr Pierre Gueriau (National Museum of Natural History, Paris), whose comments and suggestions improved the quality of the final version of the manuscript. This study was funded by the Institute of Geological Sciences of the NAS of Ukraine (research project “Fauna and flora of the Ediacaran and Palaeozoic of Ukraine: taxonomy, biostratigraphy, and biogeography”).

References

- Aisenverg D.Ye., Belenko N.G., Dedov V.S., Levenshtein M.L., Makarov I.A., Nesterenko L.P., Poletaev V.I., Popov V.S., Sokolova G.U., Fissunen O.P., & Shchegolev A.K., 1975. Stratigraficheskaya ekskursiya [Stratigraphic excursion]. [In:] Aisenverg D.Ye., Lagutina V.V., Levenshtein M.L. & Popov V.S. (Eds), *Putevoditel' ekskursii po Donetskomu basseynu* [Field excursion guidebook for the Donets Basin]. Nauka, Moscow, 201–245.
- Aisenverg D.Ye., Brazhnikova N.E., Novik K.O., Rotai A.P. & Shulga P.L., 1963. *Stratigrafiya kamennougol'nykh otlozheniy Donetskogo basseyna* [Carboniferous stratigraphy of the Donets Basin]. Publishing House of the Academy of Sciences of the Ukrainian SSR, Kyiv, 182 pp.
- Ambrose T. & Romano M., 1972. New Upper Carboniferous Chelicerata (Arthropoda) from Somerset, England. *Palaeontology* 15, 569–578.
- Anderson L.I., 1994. Xiphosurans from the Westphalian D of the Radstock Basin, Somerset Coalfield, the South Wales Coalfield and Mazon Creek, Illinois. *Proceedings of the Geologists' Association* 105, 265–275. [https://doi.org/10.1016/S0016-7878\(08\)80179-4](https://doi.org/10.1016/S0016-7878(08)80179-4)
- Anderson L.I., Dunlop J.A., Horrocks C.A., Winkelmann H.M. & Eagar R.M.C., 1997. Exceptionally preserved fossils from Bickershaw, Lancashire, UK (Upper Carboniferous, Westphalian A (Langsettian)). *Geological Journal* 32, 197–210. [https://doi.org/10.1002/\(SICI\)1099-1034\(199709\)32:3<197::AID-GJ739>3.0.CO;2-6](https://doi.org/10.1002/(SICI)1099-1034(199709)32:3<197::AID-GJ739>3.0.CO;2-6)
- Anderson L.I., Dunlop J.A., Eagar R.M.C., Horrocks C.A. & Wilson H.M., 1999. Soft-bodied fossils from the roof shales of the Wigan Four Foot coal seam, Westhoughton, Lancashire, UK. *Geological Magazine* 135, 321–329. <https://doi.org/10.1017/S0016756899002575>
- Babcock L.E. & Merriam D.F., 2000. Horseshoe crabs (Arthropoda; Xiphosurida) from the Pennsylvanian of Kansas and elsewhere. *Transactions of the Kansas Academy of Science* 103, 76–94. <https://doi.org/10.2307/3627941>
- Baily W.H., 1863. Remarks on some Coal Measures Crustacea belonging to the genus *Belinurus* König, with description of two new species from Queen's County, Ireland. *Annals and Magazine of Natural History* 11, 107–114. <https://doi.org/10.1080/00222936308681390>
- Betekhtina O.A., 1974. *Biostratigrafiya i korrelyatsiya uglenosnykh otlozheniy pozdnego paleozoya po nemorskim doustvorkam* [Non-marine bivalves, biostratigraphy and correlation of the Late Paleozoic Coal Measures]. Nauka, Novosibirsk, 178 pp.
- Bicknell R.D.C. & Pates S., 2020. Pictorial atlas of fossil and extant horseshoe crabs, with focus on Xiphosurida. *Frontiers in Earth Science* 8, 1–60. <https://doi.org/10.3389/feart.2020.00098>
- Bicknell R.D.C., Pates S. & Botton M.L., 2019. *Euproops danae* (Belinuridae) cluster confirms deep origin of gregarious behaviour in xiphosurids. *Arthropoda Selecta* 28, 549–555. <https://doi.org/10.15298/arthritis.28.4.07>
- Bicknell R.D.C., Tashman J.N., Edgecombe G.D. & Paterson J.R., 2021. Carboniferous horseshoe crab musculature suggests anatomical conservatism within Xiphosurida. *Papers in Palaeontology* 8, 1–9. <https://doi.org/10.1002/spp2.1403>
- Bölsche W., 1885. Über *Prestwichia rotundata* H. Woodw. sp. aus der Steinkohlenformation des Piesberges bei Osnabrück. *Jahresbericht des Naturwissenschaftlichen Vereins Osnabrück* 6, 268–271.
- Boyarina N.I., 2016. Makrofloristicheskiye zony srednego i verkhnego karbona (pensil'vanskoy podsystemy) Donetskogo basseyna [Macrofloristic zones of the middle and Upper Carboniferous (Pennsylvanian Subsystem) of the Donets Basin]. *Geologicheskii zhurnal* 354, 21–35. <https://doi.org/10.30836/igs.1025-6814.2016.1.97280>
- Brauckmann C., 1982. Der Schwertschwanz *Euproops* (Xiphosurida, Limulina, Euproopacea) aus dem Ober-Karbon des Piesbergs bei Osnabrück. *Osnabrücker Naturwissenschaftliche Mitteilungen* 9, 17–26.

- Bronn H.G., 1839. Fossile *Limulus*-Arten. *Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde* 1839, 489–490.
- Buckland W., 1837. The Bridgewater treatises on the power, wisdom and goodness of God as manifested in the creation. Treatise IV. Geology and mineralogy with reference to natural theology (second edition). William Pickering, London, 316 pp.
- Chernyshev B.I., 1927. Zametka o predstavityakh Xiphosura iz Donetskogo basseyna [A note on representatives of Xiphosura from the Donets Basin]. *Proceedings of the Geological Committee* 46, 645–655.
- Chernyshev B.I. 1928. Yeshche o Phyllopoda i Xiphosura Donetskogo basseyna [More on Phyllopoda and Xiphosura of the Donets Basin]. *Proceedings of the Geological Committee* 47, 519–533.
- Chernyshev B.I., 1931. Carbonicola, Anthracomya i Najadites Donetskogo basseyna [Carbonicola, Anthracomya, and Najadites of the Donets Basin]. Geological Publishing House of the Main Geological Prospecting Directorate, Moscow and Leningrad, 126 pp.
- Chernyshev B.I., 1939. Rakoobraznye [Crustaceans]. [In:] Gorsky, I.I. (Ed.), *Atlas rukovodyaschikh form iskopayemykh faun SSSR. Tom 5. Sredniy i verkhniy karbon* [Atlas of index fossils of the USSR. Volume 5. Middle and Upper Carboniferous]. Gosgeoloptekhzdat, Moscow, 141–144.
- Cockerell T.D.A., 1905. Two Carboniferous genera of xiphosurans. *American Geologist* 36, 330.
- Dernov V., 2019a. Taphonomy and paleoecology of fauna and flora from deltaic sandstones of the Mospinka Formation (middle Carboniferous) of the Donets Basin. *Geo&Bio* 18, 37–63. <https://doi.org/10.15407/gb1805>
- Dernov V.S., 2019b. K izucheniyu nemorskoï fauny mospinskoi svity (sredniy karbon, Donbass) [On the study of the non-marine fauna of the Mospino Formation (Middle Carboniferous, Donets Basin)]. *Tectonics and Stratigraphy* 46, 105–115. <https://doi.org/10.30836/igs.0375-7773.2019.208882>
- Dernov V., 2021. The earliest insect endophytic oviposition (Early Pennsylvanian, eastern Ukraine). *Visnyk of the Taras Shevchenko National University of Kyiv. Geology* 95, 16–24. <https://doi.org/10.17721/1728-2713.95.02>
- Dernov V., 2022a. Nemors'ki peletsypody mospyns'koyi svity (verkhniy bashkyr) Donetsk'koho baseynu: systematichnyy sklad, paleoekolohiya ta stratyhrafichne znachennya [Non-marine bivalves from the Mospyne Formation (upper Bashkirian) of the Donets Basin: taxonomy, paleoecology, and stratigraphic significance]. *Geologichnij zhurnal* 380, 34–56. <https://doi.org/10.30836/igs.1025-6814.2022.3.255491>
- Dernov V., 2022b. Late Bashkirian ammonoids from the Mospyne Formation of the Donets Basin, Ukraine. *Fossil Imprint* 78, 489–512. <https://doi.org/10.37520/fi.2022.021>
- Dernov V.S., 2022c. The first record of Cyclida (Pancrustacea, Multicrustacea) in the Carboniferous of the Donets Basin, Ukraine. *Collection of scientific works of the Institute of Geological Sciences of NAS of Ukraine* 15, 78–85. <https://doi.org/10.30836/igs.2522-9753.2022.267079>
- Dernov V.S., 2022d. Pryzhyttyevi poshkodzhennya cherepashok nemors'kykh dvostulkovykh molyuskiv z karbonu Donetsk'koho baseynu [Shell injuries of the Carboniferous non-marine bivalves from the Donets Basin]. *Paleontological Review* 54, 56–64.
- Dernov V., 2022e. Biostratyhafiya bashkyrs'koho yarusu (pensyl'vaniy) Donetsk'koho baseynu za faunoyu amonoidey [Ammonoid biostratigraphy of the Bashkirian (Pennsylvanian) in the Donets Basin]. *Collection of scientific works of the Institute of Geological Sciences of the NAS of Ukraine* 15, 123–137. <https://doi.org/10.30836/igs.2522-9753.2022.267080>
- Dernov V., 2023a. The first record of the trace fossils *Cochlichnus* from the Pennsylvanian continental and marine deposits in the Donets Basin, Ukraine. *Geo&Bio* 25, 109–120. <https://doi.org/10.53452/gb2508>
- Dernov V., 2023b. Horseshoe crab trace fossils *Arborichnus Romano et Meléndez*, 1985 from the Bashkirian (Carboniferous) of the Donets Basin, Ukraine. *Fossil Imprint* 79, 9–25. <https://doi.org/10.37520/fi.2023.002>
- Dernov V., 2025a. New records of Pennsylvanian phyllocarid crustaceans (Phyllocarida, Archaeostraca) from the Donets Basin, eastern Ukraine. *Geologos* 31, 1–16. <https://doi.org/10.14746/logos.2025.31.1.01>
- Dernov V.S., 2025b. A new record of the myriapod genus *Arthropleura* Jordan in Jordan et von Meyer, 1854 from the Pennsylvanian of Ukraine. *Geologichnij zhurnal* 390, 33–43. <https://doi.org/10.30836/igs.1025-6814.2025.1.301847>
- Dernov V.S. & Poletaev V.I., 2024. New geological and palaeontological data of the Dyakove Group (Carboniferous) and age-related rock formations of the central Donets Basin, Ukraine. *Geologichnij zhurnal* 386, 3–21. <https://doi.org/10.30836/igs.1025-6814.2024.1.285644>
- Dernov V.S. & Udovychenko M.I., 2019. K paleobotanicheskoy kharakteristike mospinskoy svity (sredniy karbon, Donbass) [On the palaeobotanical characteristic of the Mospino Formation]. *Visnyk of V.N. Karazin Kharkiv National University, Geology, Geography, Ecology* 51, 67–82.
- Dernov V.S. & Udovychenko N.I., 2021. Nakhodka drevneyshego konkhostraka (Crustacea: Branchiopoda) v karbone Donbassa [A record of the earliest conchostracan crustacean (Crustacea: Branchiopoda) in the Carboniferous of the Donets Basin]. *Natural Sciences: Projects, Research, Prospects. Proceedings of the 2nd International Scientific and Practical Conference (Starobilsk, 21–22 December 2021)*. Starobilsk, 31–33.
- Dewar W., 1939. *Anthraconauta* (*Anthracomya*) *minima* (Auctorum) and its associates in the Lancashire coalfield. *Bulletin of Geological Survey of Great Britain* 1, 47–66.
- Dix E. & Pringle J., 1929. On the fossil Xiphosura from the South Wales Coalfield with a note on the myriapod *Euphoberia*. *Summary of Progress, Geological Survey of Great Britain* 1928, 90–113.
- Dix E. & Pringle J., 1930. Some Coal Measure arthropods from the South Wales Coalfield. *Annals and*

- Magazine of Natural History 10, 136–144. <https://doi.org/10.1080/00222933008673194>
- Dunaeva N.M., 1969. Vidkrytyi Donbas [Open Donets Basin]. [In:] Bondarchuk, V.G. (Ed.), *Stratygraphiya URSR. Tom 5. Karbon [Stratigraphy of the Ukrainian SSR. Vol. 5. Carboniferous]*. Naukova Dumka, Kyiv, 21–48.
- Dunn D.L., 1966. New Pennsylvanian conodonts from southwestern United States. *Journal of Paleontology* 40, 1294–1303.
- Eagar R.M.C., 1981. Seasonal sedimentation in the Upper Carboniferous of England: A discussion. *The Journal of Geology* 89, 765–766. <https://doi.org/10.1086/628642>
- Eller E.R., 1938. A review of the Xiphosuran genus *Belinurus*, with the description of a new species, *B. alleghanyensis*. *Annals of the Carnegie Museum* 27, 129–150. <https://doi.org/10.5962/p.214431>
- Eros J.M., Montañez I.P., Osleger D.A., Nemyrovska T.I., Poletaev V.I., Davydov V.I. & Zhykalyak M.V., 2012. Sequence stratigraphy and onlap history of the Donets Basin, Ukraine: insight into Carboniferous icehouse dynamics. *Palaeogeography, Palaeoclimatology, Palaeoecology* 313–314, 1–25. <https://doi.org/10.1016/j.palaeo.2011.08.019>
- Feofilova A.P. & Levenstein M.L., 1963. *Osobennosti osadko- i uglenakopleniya v nizhnem i srednem karbone Donetskogo basseyna [Features of sedimentation and coal accumulation in the Lower and Middle Carboniferous of the Donets Basin]*. Publishing House of the Academy of Sciences of the USSR, Moscow, 180 pp.
- Filipiak P. & Krawczyński W., 1996. Westphalian xiphosurans (Chelicerata) from the Upper Silesia Coal Basin of Sosnowiec, Poland, *Acta Palaeontologica Polonica* 41, 413–425.
- Fisher D.C., 1977. Functional significance of spines in the Pennsylvanian horseshoe crab *Euproops danae*. *Paleobiology* 3, 175–195. <https://doi.org/10.1017/S009483730000525X>
- Fisher D.C., 1979. Evidence for subaerial activity of *Euproops danae* (Merostomata, Xiphosurida). [In:] Nitecki M.H. (Ed.), *Mazon Creek fossils*. Academic Press, New York, 379–447. <https://doi.org/10.1016/B978-0-12-519650-5.50022-9>
- Fissunen O.P., 1987. Srednekamennougol'nyye landshafty Donetskogo basseyna [Middle Carboniferous landscapes of the Donets Basin]. [In:] Bogdanova T.N. & Khozatsky L.I. (Eds.), *Palaeontology and geologic history of the palaeobasins Nedra, Leningrad*, 92–99.
- Fissunen O.P., 1991. Zonal'naya fitostratigraficheskaya shkala nizhnego i srednego karbona Donetskogo basseyna [Zonal phytostratigraphic scheme of the Lower and Middle Carboniferous of the Donets Basin]. *Geologichnij zhurnal* 3, 55–64.
- Harris R.W. & Hollingsworth R.V., 1933. New Pennsylvanian conodonts from Oklahoma. *American Journal of Science, Series 5* 25, 193–204. <https://doi.org/10.2475/ajs.s5-25.147.193>
- Haug C. Van Roy P., Leipner A., Funch P., Rudkin D.M., Schöllmann L. & Haug J.T., 2012. A holomorph approach to xiphosuran evolution – a case study on the ontogeny of *Euproops*. *Developmental Genes and Evolution* 222, 253–268. <https://doi.org/10.1007/s00427-012-0407-7>
- Higgins A.C. & Bouckaert J., 1968. Conodont stratigraphy and paleontology of the Namurian of Belgium. Brussels: Service Géologique de Belgique, *Mémoires pour servir à l'explication des cartes géologique et minières de la Belgique* 10, 1–64.
- Hind W., 1894–1896. A monograph on *Carbonicola*, *Anthracomys*, and *Naiadites*. Palaeontographical Society, London, 182 pp. <https://doi.org/10.5962/bhl.title.11877>
- Hitchcock E., 1858. Ichnology of New England; a report on the sandstone of the Connecticut Valley, especially its footmarks. W. White, Boston, 220 pp.
- Igo H. & Koike T., 1964. Carboniferous conodonts from the Omi limestone, Niigata Prefecture, central Japan (Studies of Asian conodonts, Part I). *Transactions and Proceedings of the Palaeontological Society of Japan, New Series* 53, 179–193.
- Izart A., Briand C., Vaslet D., Vachard D., Coquel R. & Maslo A., 1996. Stratigraphy and sequence stratigraphy of the Moscovian in the Donets Basin. *Tectonophysics* 268, 189–209. [https://doi.org/10.1016/S0040-1951\(96\)00224-7](https://doi.org/10.1016/S0040-1951(96)00224-7)
- Jones T.R., 1891. On some *Estheria* and *Estheria*-like shells from the Carboniferous shales of Western Scotland. *Transactions of the Geological Society of Glasgow* 9, 79–85. <https://doi.org/10.1144/transglas.9.1.79>
- Jones T.R. & Woodward H., 1899. Contributions to fossil Crustacea. *Geological Magazine* 6, 388–395. <https://doi.org/10.1017/S0016756800142487>
- Karlov N.N., 1948. *Prestwischinella tschernyschevi* sp. nov. – novyy predstavnyk Xiphosura z produktivnoyi tovshchi donets'koho karbonu [*Prestwischinella tschernyschevi* sp. nov., a new representative of Xiphosura from the Carboniferous coal-bearing succession of the Donets Basin]. *Collection of papers on palaeontology and stratigraphy*. Publishing House of the Academy of Sciences of the Ukrainian SSR; Kyiv, 41–47.
- Kozitskaya R.I., Kossenko Z.A., Lipnyagov O.M. & Nemirovskaya T.I., 1978. *Kamennougol'nyye konodonty Donetskogo basseyna [Carboniferous conodonts of the Donets Basin]*. Naukova Dumka, Kyiv, 122 pp.
- Lamsdell J.C., 2019. Evolutionary history of the dynamic horseshoe crab. [In:] Sitters H.P., Clark J.A., Smith P.A. & Niles L.J. (Eds.), *Conservation and research on shorebirds and horseshoe crabs in Delaware Bay 1997–2018. International Wader Studies* 21, 1–14.
- Lamsdell J.C., 2020. The phylogeny and systematics of Xiphosura. *PeerJ* 8: e10431. <https://doi.org/10.7717/peerj.10431>
- Lamsdell J.C., 2022. One name to rule them all: *Belinurus trilobitoides* (Buckland, 1837) is senior synonym to fourteen named species. *Journal of Paleontology* 96, 237–241. <https://doi.org/10.1017/jpa.2021.84>
- Lamsdell J.C. & Clapham M.E., 2021. *Belinurus* Bronn, 1839 (Chelicerata, Xiphosura) has priority over *Bellinurus* Pictet, 1846. *Journal of Paleontology* 95, 1352–1355. <https://doi.org/10.1017/jpa.2021.53>
- Latreille P.A., 1802. Histoire naturelle générale et particulière des crustacés et des insectes: ouvrage faisant suite

- aux Oeuvres de Leclerc de Buffon, et partie du Cours complet d'histoire naturelle rédigé par C.S. Sonnini, 7. F. Dufart, Paris, 413 pp. <https://doi.org/10.5962/bhl.title.15764>
- Linnaeus C., 1758. *Systema Naturae*. Vol. 1. 10th edition. Laurentius Salvius, Holmiae, 824 pp.
- Leach W., 1819. Entomostracés. [In:] Levrault F. (Ed.), *Dictionnaires des Sciences Naturelles*, 14. Strasbourg, pp. 524–543.
- Logvinenko N.V., 1953. *Litologiya i paleogeografiya produktivnoy tolshchi donetskogo karbona* [Lithology and palaeogeography of the Carboniferous coal-bearing deposits of the Donets Basin]. Publishing House of the Kharkiv University, Kharkiv, 436 pp.
- Lucas S.G., Lerner A.J., Dimichele W.A., Cantrell A.K., Suazo T.L. & Chaney D.S., 2014. Xiphosurid fossils from the Pennsylvanian Beeman Formation, Otero County, New Mexico. *New Mexico Geological Society Guidebook*, 65th Field Conference, *Geology of the Sacramento Mountains Region*, 311–314. <https://doi.org/10.56577/FFC-65.311>
- Lustri L., Laibl L. & Bicknell R.D.C., 2020. A revision of *Prolimulus woodwardi* Fritsch, 1899 with comparison to other highly paedomorphic belinurids. *PeerJ* 9, e10980. <https://doi.org/10.7717/peerj.10980>
- Meek E.B., 1867. Note on a new genus of fossil Crustacea. *The American Journal of Science and Arts (Series 2)* 43, 394–395.
- Meek F.B. & Worthen A.H., 1865. Notice of some new types of organic remains, from the Coal Measures of Illinois. *Proceedings of the Academy of Natural Sciences* 12, 41–45.
- Mikulic D.G., 1998. Xiphosura. [In:] Shabica C.W. & Hays A.A. (Eds.), *Richardson's guide to the fossil fauna of Mazon Creek*. Northeastern Illinois University Press, Chicago, 134–139.
- Müller O., 1785. Entomostraca, seu, Insecta testacea quae in aquis Daniae et Norvegie reperit, descripsit et iconibus illustravit. F.W. Thiele, Hauniae, 135 pp. <https://doi.org/10.5962/bhl.title.14414>
- Mychko E.V. & Dernov V.S., 2019. Novyye nakhodki trilobitov v srednem karbone Donetskogo basseyna [New record of trilobites in the middle Carboniferous of the Donets Basin]. *Bulletin of the Moscow Society of Naturalists (Geological Section)* 94, 52–63.
- Nemyrovska T.I., 1999. Bashkirian conodonts of the Donets Basin, Ukraine. *Scripta Geologica* 119, 1–116.
- Nemyrovska T.I., 2017. Late Mississippian–Middle Pennsylvanian conodont zonation of Ukraine. *Stratigraphy* 14, 299–318. <https://doi.org/10.29041/strat.14.1-4.299-318>
- Nemyrovska T.I., & Yefimenko V.I., 2013. Seredniy karbon (nyzniy pensyl'vaniy) [Middle Carboniferous (Lower Pennsylvanian)]. [In:] Gozhyk P.F. (Ed.), *Stratyhafiya verkh'n'oho proterozoyu ta fanerorozoyu Ukrayiny. Tom 1. Stratyhafiya verkh'n'oho proterozoyu, paleozoyu ta mezozoyu Ukrayiny* [Stratigraphy of the Upper Proterozoic and Phanerozoic of Ukraine. Vol. 1. Stratigraphy of the Upper Proterozoic, Palaeozoic and Mesozoic]. LAT&K, Kyiv, 283–303.
- Newberry J.S., 1856. Description of several new genera and species of fossil fishes, from the Carboniferous strata of Ohio. *Proceedings of the Academy of Natural Sciences of Philadelphia* 8, 96–100.
- Novik E.O., 1974. *Zakonomernosti razvitiya kamennougol'noy flory yuga Yevropeyskoy chasti SSSR* [Development of the Carboniferous flora of the south part of the European part of the USSR]. Naukova Dumka, Kyiv, 139 pp.
- Packard A.S., 1886. On the Carboniferous xiphosurous fauna of North America. *Memoirs of the National Academy of Sciences* 3, 143–157. <https://doi.org/10.5962/bhl.title.14755>
- Pictet F.J., 1854. *Traite élémentaire de paléontologie*. Vol. 2. Baillière, Paris, 727 pp.
- Poletaev V.I. & Vdovenko M.V., 2013. Nyzhniy karbon (Missisipi) [Lower Carboniferous (Mississippian)]. [In:] Gozhik P.F. (Ed.), *Stratyhafiya verkhnyogo proterozoyu i fanerorozoyu Ukrainy. Tom 1. Stratyhafiya verkhnyogo proterozoyu, paleozoyu i mezozoyu* [Stratigraphy of the Upper Proterozoic and Phanerozoic of Ukraine. Volume 1. Stratigraphy of the Upper Proterozoic, Palaeozoic and Mesozoic]. LAT&K, Kyiv, 250–283.
- Poletaev V.I., Vdovenko M.V., Shchoholev O.K., Boyarina N.I., & Makarov I.A., 2011. Stratotypy rehional'nykh stratyhrafichnykh pidrozdiliv karbonu i nyzhn'oyi permi Dono-Dniprovs'koho prohynu [Stratotypes of the Carboniferous and Lower Permian regional stratigraphic units of the Don-Dnipro Downwarp]. Logos, Kyiv, 234 pp.
- Popov A.V., 1979. Kamennougol'nyye ammonoidei Donbassa i ikh stratigraficheskoye znachenije [Carboniferous ammonoids of the Donets Basin and their stratigraphic significance]. *Transactions of the All-Union Geological Institute, New series* 220, 1–119.
- Posudiyevsky A.B., 1977. K izucheniyu fatsial'nogo sostava porod neposredstvennoy krovli ugol'nykh plastov v Donbasse [On the study of depositional environments of the roof shales of the coal beds in the Donets Basin]. *Lithology and mineral resources* 3, 156–159.
- Prantl F. & Příbyl A., 1955. Ostroropi (Xiphosura) československého karbonu. *Sborník Ústředního ústavu geologického* 22, 379–424.
- Pruvost P., 1930. La faune continentale du terrain houiller de la Belgique. *Mémoires du Musée royal d'histoire naturelle de Belgique* 44, 103–282.
- Reznikov A.I., 1993. D'yakovskaya seriya i yeye polozheniye v stratigraficheskoy skheme Donbassa [The Dyakove Group and its stratigraphic position in the Carboniferous succession of the Donets Basin]. *Geologicheskij zhurnal* 262, 52–57.
- Roemer F., 1883. Über eine Art der limuliden-gattung *Belinurus* aus dem Steinkohlengebirge Oberschlesiens. *Zeitschrift der Deutschen Geologischen Gesellschaft* 35, 429–432.
- Sachsenhofer R.F., Privalov V.A., & Panova E.A., 2012. Basin evolution and coal geology of the Donets Basin (Ukraine, Russia): An overview. *International Journal of Coal Geology* 89, 26–40. <https://doi.org/10.1016/j.coal.2011.05.002>
- Schultka S., 1994. *Bellinurus* cf. *truernanii* (Merostomata) aus dem tiefen Oberkarbon (Namur B/C) von Frodenberg (Nordrhein-Westfalen, Deutschland).

- Paläontologische Zeitschrift 68, 339–349. <https://doi.org/10.1007/BF02991347>
- Schwarzbach M., 1962. Die Merostomata aus dem niederrheinisch-westfälischen Oberkarbon. *Das Karbon der subvariscischen Saumsenke. Ein Symposium. Teil 2. Fortschritte in der Geologie von Rheinland und Westfalen* 3, 803–818.
- Selden P.A., Simonetto L. & Marsiglio G., 2019. An effaced horseshoe crab (Arthropoda: Chelicerata: Xiphosura) from the Upper Carboniferous of the Carnic Alps (Friuli, NE Italy). *Rivista Italiana di Paleontologia e Stratigrafia* 125, 333–342.
- Shpinev E.S., 2018. New data on Carboniferous xiphosurans (Xiphosura, Chelicerata) of the Donets Coal Basin. *Paleontological Journal* 52, 271–283. <https://doi.org/10.1134/S0031030118030127>
- Shulga V.F., Zdanowski A., Zaitseva L.B., Ivanova A.V., Ivanina A.V., Korol N.D., Kotasova A., Kotas A., Kostyk I.E., Lelyk B.I., Miger T., Manichev V.I., Matrofailo M.N., Ptak B., Savchuk V.S., Sedaeva G.M. & Stepanenko Ya.G., 2007. *Korrelyatsiya karbonovykh ughlenosnykh formatsiy L'vovsko-Volynskogo i Lyublinskogo basseyinov* [Correlation of the Carboniferous coal-bearing formations of the Lviv-Volyn and Lublin basins]. Varta, Kyiv, 427 pp.
- Siegfried P., 1972. Ein Schwertschwanz (Merostomata, Xiphosurida) aus dem Oberkarbon von Ibbenbüren/Westf. *Paläontologische Zeitschrift* 46, 180–185. <https://doi.org/10.1007/BF02990151>
- Stainier X., 1938. *Belinurus* et *Vetacapsula* nouveaux du houiller Belge. *Bulletin du Musée royal d'Histoire naturelle de Belgique* 14, 1–13.
- Tashman J.N., Feldmann R.M. & Schweitzer C.E., 2019. Morphological variation in the Pennsylvanian horseshoe crab *Euproops danae* (Meek & Worthen, 1865) (Xiphosurida, Euproopidae) from the lower Mercer Shale, Windber, Pennsylvania, USA. *Journal of Crustacean Biology* 39, 396–406. <https://doi.org/10.1093/jcbiol/ruz030>
- Trueman A.E. & Weir J., 1948. A monograph of British Carboniferous non-marine Lamellibranchia. Part IV. *Monographs of the Palaeontographical Society* 102, 75–100. <https://doi.org/10.1080/02693445.1948.12035642>
- Ustinovsky Yu.B., 1993. Kayal'skiy etap (pozdnii bashkir) [Kayalian stage (Late Bashkirian)]. [In:] Tsegelnyuk P.D. (Ed.), *Geologicheskaya istoriya territorii Ukrainy* [Geological history of Ukraine]. Naukova Dumka, Kyiv, 141–144.
- Van Hinsbergen D.J.J., Abels H.A., Bosch W., Boekhout F., Kitchka A., Hamers M., van der Meer D.G., Geluk M. & Stephenson R.A., 2015. Sedimentary geology of the middle Carboniferous of the Donbas region (Dniepr-Donets basin, Ukraine). *Scientific Reports* 5 (9099). <https://doi.org/10.1038/srep09099>
- Warth M., 1967. Die nichtmarinen Muscheln des Westfal A unter besonderer Berücksichtigung des Ruhrkarbons und einige grundlegende Erkenntnisse zur Taxonomie. Westdeutscher Verlag, Köln, 125 s.
- Weber V.N., 1933. Trilobity Donetskogo basseyina [Trilobites of the Donets Basin]. *Transactions of the All-Union Geological Exploration Department of NKTP of the USSR* 225, 1–95.
- Wignall P.B. & Sims M.J., 1990. Pseudoplankton. *Palaeontology* 33, 359–378.
- Williamson W.C., 1849. On the microscopic structure of the scales and dermal teeth of some ganoid and placoid fish. *Philosophical Transactions of the Royal Society of London* 139, 435–474. <https://doi.org/10.1098/rstl.1849.0023>
- Woodward H., 1867. On some points in the structure of the Xiphosura having reference to their relationship with the Eurypteridae. *Quarterly Journal of the Geological Society of London* 23, 28–37. <https://doi.org/10.1144/GSL.JGS.1867.023.01-02.11>
- Woodward H., 1893. Note on a new British species of *Cyclus* from the Coal Measures of Bacup, Lancashire. *Geological Magazine* 10, 28–29.
- Woodward H., 1918. Fossil arthropods from the Carboniferous rocks of Cape Breton, Nova Scotia; and from the Upper Coal Measures, Sunderland, England. *Geological Magazine* 5, 462–471. <https://doi.org/10.1017/S0016756800203944>
- Woodward H. & Etheridge R., 1873. Explanation of sheet 23. *Memoirs of the Geologic Survey, Scotland*, 1–99.
- Wright W.B., 1934. The fresh-water fauna of the Lower Measures of Lancashire. *Summary of progress of the Geological Survey of Great Britain and the Museum of Practical Geology for the year 1933* 2, 8–23.
- Yefimov I.N., 1934. Raspredeleniye organicheskikh ostatkov v krovle plastov kamennykh ugley Donetskogo basseyina [Distribution of fossils in the roof shales of the coal beds in the Donets Basin]. *Reports of the Academy of Sciences of the USSR* 6, 123–125.
- Zalessky M.D., 1907. Materialy po kamennougol'noy flore Donetskogo basseyina. I. Rastitel'nyye ostatki kollektsii V. Domgera [Materials on the Carboniferous flora of the Donets Basin. I. Plant remains of V. Domher's collection]. *Proceedings of the Geological Committee* 26, 351–422.
- Zaritskiy P.V., 1970. *Mineralogiya i geokhimiya diageniza ughlenosnykh otlozheniy (na primere Donetskogo basseyina)*. Chast' 1. [Mineralogy and geochemistry of diagenesis of coal-bearing deposits (based on materials from the Donets Basin). Part 1]. Kharkiv University Press, Kharkiv, 225 pp.
- Zhemchuzhnikov Yu.A., Yablokov V.S., Bogolyubova L.I., Botvinkina L.N., Feofilova A.P., Ritenberg M.I., Timofeev P.P. & Timofeeva Z.V., 1960. Stroyeniye i usloviya nakopleniya glavnykh ughlenosnykh svit i ugol'nykh plastov srednego karbona v Donetskom basseyne. Chast' 2 [Structure and depositional conditions of the main coal-bearing formations and coal seams of the middle Carboniferous in the Donets Basin. Part 2]. *Transactions of the Geological Institute* 15, 1–247.
- Zittel K.A. von & Eastman C.R., 1913. Textbook of Palaeontology. Macmillan, London, 839 pp. <https://doi.org/10.5962/bhl.title.2100>